

G protein-mediated stretch reception

Ursula Storch,¹ Michael Mederos y Schnitzler,¹ and Thomas Gudermann^{1,2}

¹Walther-Straub-Institute of Pharmacology and Toxicology, University of Munich, and ²Munich Heart Alliance, Munich, Germany

Submitted 16 August 2011; accepted in final form 3 January 2012

Storch U, Mederos y Schnitzler M, Gudermann T. G protein-mediated stretch reception. *Am J Physiol Heart Circ Physiol* 302: H1241–H1249, 2012. First published January 6, 2012; doi:10.1152/ajpheart.00818.2011.—Mechanosensation and -transduction are important for physiological processes like the senses of touch, hearing, and balance. The mechanisms underlying the translation of mechanical stimuli into biochemical information by activating various signaling pathways play a fundamental role in physiology and pathophysiology but are only poorly understood. Recently, G protein-coupled receptors (GPCRs), which are essential for the conversion of light, olfactory and gustatory stimuli, as well as of primary messengers like hormones and neurotransmitters into cellular signals and which play distinct roles in inflammation, cell growth, and differentiation, have emerged as potential mechanosensors. The first candidate for a mechanosensitive GPCR was the angiotensin-II type-1 (AT₁) receptor. Agonist-independent mechanical receptor activation of AT₁ receptors induces an active receptor conformation that appears to differ from agonist-induced receptor conformations and entails the activation of G proteins. Mechanically induced AT₁ receptor activation plays an important role for myogenic vasoconstriction and for the initiation of cardiac hypertrophy. A growing body of evidence suggests that other GPCRs are involved in mechanosensation as well. These findings highlight physiologically relevant, ligand-independent functions of GPCRs and add yet another facet to the polymodal activation spectrum of this ubiquitous protein family.

mechanosensation; G protein-coupled receptor; angiotensin-II type-1 receptor; agonist independent

THIS ARTICLE is part of a collection on **G Protein Kinase A Signaling in Cardiovascular Physiology and Disease**. Other articles appearing in this collection, as well as a full archive of all collections, can be found online at <http://ajpheart.physiology.org/>.

Introduction

Cells have the ability to sense numerous chemical and physical stimuli, such as hormones and neurotransmitters, but also pain, electric fields, photons, temperature, and mechanical deformations. The capability of cells to sense physical forces and convert them into electrical and biochemical signals is termed “mechanosensation” and comprises mechanoreception as well as mechanotransduction. Mechanosensation is important for the senses of hearing, touch, and balance and for proprioception and visceroreception. Mechanosensitive cells exhibit mechanosensors, e.g., in nerve endings of the skin, in skeletal muscle, and in sensory hair cells of the inner ear (13). Mechanosensation also plays a role in the regulation of blood vessel diameter. In the endothelium as well as in smooth muscle cells of small resistance arteries, mechanosensitive structures are present and harbor key molecular components

for the autoregulation of blood flow in these tissues (76) without neuronal involvement. Although mechanosensitive cells are often well characterized, the identity of mechanosensitive subcellular structures and proteins is still widely unknown. A prominent example of as yet unknown molecular correlates underlying mechanosensation is represented by myogenic vasoconstriction, also known as the Bayliss effect: smooth muscle cells of small resistance arteries are mechanosensitive and have the intrinsic property to constrict in response to elevated intraluminal pressure (20). Such an inherent vascular regulatory mechanism allows for the functional plasticity of the cardiovascular system and is of prime physiological relevance, because it makes a significant contribution to basal vascular tone and peripheral vascular resistance as well as capillary hydrostatic pressure and organ perfusion. This review provides an overview of mechanosensitive proteins with a particular focus on the vascular system.

How Can Mechanical Forces Act on Mechanosensitive Integral Membrane Proteins?

It is commonly accepted that mechanical forces act on cell membranes by increasing the membrane tension, thereby influencing integral membrane proteins. Increased membrane tension is the key driving force (80) and can be observed for instance as membrane stretch. To date, two main concepts are discussed as to how integral membrane proteins might perceive mechanical force (49). First, the “tethered” model, which is

Address for reprint requests and other correspondence: M. Mederos y Schnitzler, Walther-Straub-Inst. of Pharmacology and Toxicology, Univ. of Munich, Goethestr. 33, 80336 Munich, Germany (e-mail: mederos@lrz.uni-muenchen.de).

based on the assumption that intra- or/and extracellular anchorage to other proteins acts as a molecular spring that confers mechanosensitivity to integral membrane proteins. The second idea is the “membrane” model, which supposes that altered membrane tension changes the lateral pressure profile (10) at the protein/phospholipid bilayer boundary to provoke conformational changes of integral membrane proteins.

At present, it is still unknown which model would apply best to myogenic vasoconstriction and how membrane stretch and membrane tension are maintained over quite a long time period during myogenic vasoconstriction. An increase in intraluminal pressure initially results in higher wall tension, thereby inducing myogenic vasoconstriction which in turn decreases wall tension (77). Mechanistically, the relationship between reduced wall tension and maintenance of myogenic tone is not understood. Furthermore, it is not clear whether wall tension of an intact blood vessel is equivalent to membrane tension of isolated smooth muscle cells. Moreover, it would appear rather unlikely that only one specific mechanosensitive protein is responsible for myogenic vasoconstriction rather than several different proteins that synergistically operate in the initiation, maintenance, and termination of myogenic tone.

Criteria for Mechanosensitive Integral Membrane Proteins

Because proteins involved in mechanically induced signaling cascades are often indiscriminately termed mechanosensitive, although they do not directly sense membrane stretch, some specific criteria should be established to unambiguously classify directly mechanosensitive proteins (14, 61). For inherently mechanosensitive proteins, there should be a strict and obvious correlation between the amplitude of the mechanical stimulus and protein activation, e.g., single channel activity of ion channels, speed of the enzymatic reaction of membrane-bound enzymes, and G protein activation of G protein-coupled receptors (GPCRs). Furthermore, the strength of the mechanical stimulus should be reflected by the magnitude of protein activation. For example, a more intense mechanical force should cause a faster protein response. To this end, a force coefficient analogous to the Q_{10} value describing the activation rate induced by a 10°C increase in temperature should be established to characterize bona fide mechanosensitive proteins (61). A high force coefficient would indicate pronounced mechanosensitivity. In addition, mechanically induced direct protein activation must occur within the protein's typical reaction time, indicating that the mechanical stimulus is adequate and physiologically meaningful so that the activation time of mechanically stimulated proteins should be very short, that is, <5 ms for ion channels (37), <50 μ s for membrane-bound enzymes (100), and <500 ms for G protein activation by GPCRs (97). Furthermore, it should be possible to suppress mechanically induced protein activation by channel blockers, inhibitors, or inverse agonists. As a more direct proof of concept, mechanically induced protein activation should result in detectable conformational changes. As the ultimate criterion for intrinsic mechanosensitivity, purified proteins should be inserted into an artificial bilayer and should still be mechanosensitive (14, 61), which was convincingly demonstrated for bacterial mechanosensitive channels (91). However, to date it is frequently difficult to discriminate between direct and indirect mechanosensitivity due to limitations of the technical

resolution and the experimental designs chosen. However, it should be kept in mind that mechanosensory protein complexes may consist of several distinct components that are neatly orchestrated to sense and transduce mechanical signals (9). By this means, a given protein may not be mechanosensitive *eo ipso* but may still react to mechanical cues in conjunction with other proteins within macromolecular complexes. It still remains elusive whether the formation of such protein complexes simply enhances a weak inherent mechanosensitivity of single protein subunits or whether it is the macromolecular assembly as such that accounts for novel mechanosensing qualities.

Potential Mechanosensors Discussed So Far

Until now, a variety of different proteins has been suggested as direct mechanosensors (see Fig. 1). For example, intracellular structures such as dense bands are discussed as mechanosensors (54). In arterioles these membrane-associated proteins consisting of vinculin and α -actinin (88) are enriched at the abluminal aspect of the blood vessel (32). This particular localization is compatible with a sensory function of these proteins. However, dense bands may serve rather as stabilizers of vascular smooth muscle cells. Other intracellular structural proteins like desmin are not involved in myogenic vasoconstriction (55). However, they may determine the rigidity of smooth muscle cells and thus of the vascular wall (50). Furthermore, the actin cytoskeleton might play a role for mechanosensation by establishing a physical link between mechanosensitive proteins (47). There is no doubt that the actin cytoskeleton influences myogenic vasoconstriction (15, 16, 31), with short- and long-term adaption of the cytoskeleton reflecting the plasticity of small resistance arteries in response to different strains (69).

Whereas the cytoskeleton may act as a tether connecting mechanosensitive proteins, the extracellular matrix may serve as an anchor relaying external relative movements to integral membrane proteins (39). This kind of mechanosensitive protein complexes is best understood in *Caenorhabditis elegans* touch receptor neurons (9, 28).

Integrins are heterodimeric glycoproteins in the plasma membrane linking components of the extracellular matrix like fibronectin, laminin, and collagen with intracellular structural proteins like dense bands. Especially with regard to myogenic vasoconstriction, integrins are discussed as mechanosensors (19, 38, 59, 64) because of their special role as linkers of intra- and extracellular proteins (40). As integrins impact the actin cytoskeleton and vice versa (92), it is still unsettled whether they primarily act as direct mechanosensors or whether they are involved in mechanotransduction as long-term modulators. As an additional facet of cell regulation, integrins impact ion channel function (101, 108, 109), thus further increasing the difficulty to precisely define their physiological role in mechanosensation.

Another family of proteins, tyrosine kinases of the Src family, was also touted as mechanosensors (84) or mechanotransducers (103, 107) because of their impact on the actin cytoskeleton via integrins (30). However, in HEK293 cells neither endogenous nor overexpressed receptor tyrosine kinases were mechanosensitive (62), leaving the role of tyrosine kinases in mechanosensation ill-defined.




Fig. 1. Possible mechanosensors. Various proteins that have been discussed as potential mechanosensors in different cells are displayed. These proteins have been proposed to be activated by membrane stretch or by fluid shear stress. Possible mechanosensors are highlighted in color and nonmechanosensitive proteins in gray. PLC β , phospholipase C β ; PLA $_2$, phospholipase A $_2$; AA, arachidonic acid; AC, adenylyl cyclase; PIP $_2$, phosphatidylinositol 4,5-bisphosphate; IP $_3$, inositol trisphosphate; DAG, diacylglycerol; Src, Src tyrosine kinase; cAMP, adenosine 3',5'-cyclic monophosphate; ATP, adenosine-5'-triphosphate; Piezo, piezo1 and piezo 2, a new family of mechanosensitive cation channels; TRP, transient receptor potential; TREK-1, TWIK-related potassium channel; Deg, degenerin; ENaC, epithelial sodium channel.

Two membrane-associated enzymes, phospholipase A $_2$ (PLA $_2$) and phospholipase C (PLC), have been convincingly implicated in mechanosensation. Slight changes of osmolarity entailed enhanced PLA $_2$ enzymatic activity (53), and direct mechanical stimulation of skeletal muscle cells potentiated PLA $_2$ but not PLC activity (94). On the contrary, PLC could be mechanically activated in mesangial cells (46). Especially in blood vessels, mechanical activation of PLC was observed (43, 60, 70, 78). While PLC can be classified as indirectly mechanosensitive (62), it is not clear at present whether PLA $_2$ reacts directly or indirectly to mechanical stimuli (41). However, there is strong experimental support of the notion that both proteins play a key role in mechanotransduction.

Potentially Mechanosensitive Ion Channels

In the past, various ion channels have been implicated as molecular components of a mechanotransduction pathway. With regard to myogenic vasoconstriction, chloride channels (23, 36, 72, 110) and cation channels (105, 106) were considered as mechanosensors. Furthermore, candidate channels of the degenerin (Deg)/epithelial sodium channel (ENaC)/acid-sensing ion channel family were suggested to be endowed with mechanosensing properties in specialized protein complexes (24, 25, 95) compared with mechanosensitive complexes in *C. elegans*. Notably, ENaC channels were particularly receptive to shear stress when endogenously (102) or heterologously expressed (11, 87) even after reconstitution in an artificial bilayer (4), pointing to their inherent mechanosensitivity.

Lately, several members of the transient receptor potential (TRP) family of cation channels (17, 65), especially TRPC1, TRPC6, TRPM4, TRPM7, TRPV2, TRPV4, and TRPP1 (PKD2) (nomenclature according to the International Union of Pharmacology), have attracted considerable attention as poten-

tially mechanosensitive ion channels (42, 85, 113). While TRPC1 was primarily suggested to be the endogenous mechanosensitive ion channel in *Xenopus* oocytes (58), this concept has been recently challenged (22, 35).

TRPC6 and TRPM4 were also regarded as direct mechanosensors. The latter two proteins are depolarizing cation channels involved in the constriction of resistance arteries as evidenced by a profound suppression of myogenic vasoconstriction of cerebral arteries after preincubation with specific antisense oligonucleotides (27, 105). In accord with the role of TRPC6 for myogenic responsiveness, inherent mechanosensitivity of TRPC6 channels was reported (89) but could not be verified later on (35, 41, 62). TRPM4 channel activity is tightly regulated by free intracellular calcium (73) and protein kinase C (26), and in a heterologous overexpression system, TRPM4 was also activated to membrane stretch (66), and clarification of the various mechanisms by which TRPM4 channel function is regulated will require additional studies. However, the channel protein's inherent sensitivity to membrane stretch is still a moot issue (96). Likewise, the activation of TRPM7 by mechanical stimuli has been reported to contribute to regulatory volume changes in epithelial cells (74, 75) but was subsequently called into question after thorough electrophysiological analysis of the overexpressed protein (7). There is hardly any solid experimental evidence that TRPM7 is in any traditional sense a mechanosensitive ion channel. Probably, the dilution of intracellular magnesium or magnesium nucleotides caused by hypotonic cell swelling rather than membrane stretch may have caused TRPM7 activation when investigating regulatory volume changes in epithelia (74, 75). Thus results obtained solely by hypotonic cell swelling as a mechanical stimulus have to be interpreted with great caution.

Thermosensitive TRPV2 channels may also serve as mechanosensors in vascular smooth muscle and may exhibit inherent mechanosensitivity upon functional expression of recombinant proteins (68). With regard to TRPV4, there is a considerable body of experimental evidence to support the notion that these ion channels are not directly gated by mechanical forces but are activated downstream of a signaling pathway, encompassing PLA₂ and cytochrome-*P450* epoxygenase (99, 104) with PLA₂, possibly representing the cellular mechanosensor. Furthermore, Deg/ENaC rather than TRPV channels were recently identified as major mechanosensors in *C. elegans* nociceptor neurons (34), whereas TRPV channels are supposed to be activated downstream of Deg/ENaC channels.

Whereas homomeric TRPP1 (PKD2) channels encoded by polycystic kidney disease 2 gene (PKD2) do not appear to be directly mechanosensitive (48), TRPP1 (PKD2) proteins complexed with TRPV4 or PKD1 (previously known as polycystin-1) might represent crucial constituents of mechanosensitive protein complexes in the vasculature (48, 86). Heteromeric TRPP1 (PKD2)/TRPV4 protein complexes have been advocated as polymodal sensory complexes that mediate flow-sensitive cation influx (48). Similarly, TRPP1 (PKD2)/PKD1 complexes in the primary cilium of kidney cells were reported to mediate mechanosensation (71), whereas in vascular smooth muscle cells, TRPP1 (PKD2) does not appear to be directly mechanosensitive but may rather inhibit endogenous stretch-activated cation channels, an effect which is abrogated by interaction with PKD1 (86). In addition, TRPP1 (PKD2) proteins are supposed to associate with TRPC1 (93). However, physiological corollaries of these TRPP1 (PKD2)/TRPC1 complexes remain elusive. Thus the proposed mechanosensitivity of TRPP1 (PKD2) is far from being understood mechanistically and presently lacks conceptual consistency.

Based on the aforementioned criteria for direct mechanosensitivity, no TRP channel can be classified as a direct mechanosensor up to now. However, there is circumstantial evidence to suggest that TRP channels play a modulatory role in mechanotransduction as crucial downstream effector proteins. However, there is compelling evidence of direct mechanosensitivity of a different cation channel, TREK-1, a two pore-domain background potassium channel. Membrane stretch activates TREK-1 channels with half-maximal activation at about -40 mmHg of negative pressure (56, 79). The mechanosensitivity of TREK-1 is independent of the expression model, the technical configuration, and the type of mechanical stimulation. Furthermore, TREK-1 gene-deficient mice exhibit reduced touch sensitivity (2), providing *in vivo* evidence for neuronal mechanosensation. Although TREK-1 was not involved in myogenic vasoconstriction in mesenteric arteries, TREK-1 plays a role for pressure-induced vasodilatation in the skin (33), highlighting distinct mechanosensory mechanisms in different tissues.

Recently, two new proteins, *piezo1* and *piezo2* (also known as *Fam38a* and *Fam38b*), were described as a new class of mechanosensitive ion channels (18). These integral membrane proteins are characterized by 24 to 36 predicted transmembrane segments and were found to be expressed in various tissues like bladder, colon, and lung where they may represent molecular correlates of mechanosensitive cation channels. Furthermore, *piezo2*, which is highly expressed in dorsal root ganglia, was found to play an important role for mechanosensation in these

somatosensory neurons (15). In accord with these observations, the amphiphilic peptide GsMTx4 was found to inhibit mechanically induced *piezo1* currents upon heterologous expression (5). The latter peptide is an inhibitor of mechanosensitive ion channels, yet does not appear to block these channels directly, but acts by insertion into the membrane bilayer, thus modifying the lipid composition surrounding integral membrane proteins, resulting in an altered lateral pressure profile at the phospholipid/protein interface (90). Thus future studies will have to show whether *piezo* proteins are in fact pore forming subunits of mechanosensitive ion channels or indispensable accessory subunits of a larger mechanosensitive protein complex.

Interestingly, voltage-gated channels appear to be mechanosensitive as well (51, 52, 67). This property may impact the regulation of heart beat, since GsMTx4 was shown to inhibit atrial fibrillation (8).

GPCRs as Mechanosensors

Recently, GPCRs, generally known as mediators of light, olfactory and gustatory stimuli, as well as of the vast majority of hormones and neurotransmitters, were appreciated as mechanosensors. GPCRs exhibit distinct functions in many physiological processes like inflammation, cell growth, and differentiation. Interestingly, the G_{q/11} protein-coupled angiotensin-II type-1 receptor (AT₁R), a cardinal component of the renin-angiotensin regulatory system, was identified as the first mechanosensitive GPCR (115). Instead of autocrine angiotensin-II release (83), Zou et al. (115) demonstrated that agonist-independent mechanical AT₁R activation is one of the main mechanisms mediating stretch-induced hypertrophy of cardiomyocytes. Stretch-induced receptor activation could be completely prevented by deploying specific inverse AT₁R agonists (111, 115). Involvement of endogenous angiotensin II was ruled out using angiotensinogen-deficient mice and a neutralizing antibody directed against angiotensin II.

Apart from the involvement of AT₁R in the pathophysiological state of cardiac hypertrophy, AT₁R were identified as mechanosensors participating in the physiological process of myogenic vasoconstriction (62). Membrane stretch in a physiological range was shown to activate G_{q/11} protein-coupled receptors independently of endogenous agonist (62), demonstrated firstly by employing an angiotensin-converting enzyme inhibitor to suppress endogenous angiotensin II secretion and secondly by the fact that in rat cerebral arteries, the endothelium does not express chymases which generate angiotensin II and which are present in other species (44, 63). In addition, a neutralizing antibody directed against angiotensin II did not affect mechanically induced AT₁R signaling (62). Furthermore, in the conditioned medium of mechanically stimulated myogenic arteries, angiotensin II was not detectable by radioimmunoassay and a sensitive bioassay, thus ruling out any contribution of endogenous angiotensin II on mechanical receptor activation (62). However, under physiological conditions, basal angiotensin II levels are likely to sensitize the receptor's response toward mechanical stimulation.

When compared with Zou et al. (115) in analyzing very distal steps of the signaling cascade induced by receptor activation, Mederos y Schnitzler and colleagues focused on more proximal events. In the latter study mechanically induced receptor activation was PLC dependent, which is in line with

former studies showing that PLC isoforms are important elements of signaling pathways involved in mechanotransduction in myogenic arteries (43, 60, 70, 78). Stretch-induced receptor activation leads to GTP loading of $G_{q/11}$ proteins, PLC activation, and subsequently diacylglycerol-mediated TRPC channel activation, allowing for sodium and calcium influx. Beyond PLC, PLA_2 might also act as mechanotransducers potentiating TRPC6 activity by 20-hydroxyeicosatetraenoic acid production in concert with ω -hydroxylase (41). However, it is still not clear whether PLA_2 isoforms are intrinsically mechanosensitive, whereas in contrast PLCs are most probably indirectly mechanosensitive. Altogether, these findings revise the present views of GPCR function in general and they shed new light on the mechanism of myogenic vasoconstriction. In contrast to former concepts of mechanosensation, GPCRs are principally mechanosensitive and TRPC channel activation occurs downstream of mechanically induced GPCR activation, indicating that TRPC channels are only indirectly mechanosensitive. In smooth muscle cells TRPC channel activation leads to depolarization of the membrane potential, thus increasing the channel open probability of L-type voltage-gated calcium channels resulting in calcium influx, elevation of the intracellular calcium concentration, force development, and smooth muscle cell contraction (62, 98).

Mechanical GPCR activation is not restricted to AT_1R s. Although the $G_{q/11}$ -coupled endothelin type-1a (ET_{1A}) receptors were initially described as insensitive to mechanical stimulation pointing to a rather exceptional property of AT_1R s as mechanosensors (115), Mederos y Schnitzler et al. showed that a large variety of $G_{q/11}$ protein-coupled receptors, including the endothelin ET_{1A} receptor, are directly mechanosensitive (62). Furthermore, the propensity to react to mechanical stimuli varied from the histamine H_1 receptor, exhibiting the strongest cell responses upon mechanical stimulation, followed by the AT_1R , the muscarinic M_5 receptor, and finally the vasopressin V_{1A} receptor. As yet, the structural basis for such differential mechanosensitivity is unknown.

It is reasonable to propose that the mechanosensitivity of $G_{q/11}$ protein-coupled receptors is involved in other physiological processes as well and may even be a general property of most, if not all, $G_{q/11}$ protein-coupled receptors. In smooth muscle cells, mechanosensitivity depends on receptor density (62), indicating that the specificity of cells, tissues, and organs to be activated by mechanical stimulation is regulated by the expression levels of the GPCRs.

As mentioned before, directly mechanosensitive proteins are characterized by mechanically induced conformational changes. In line with this prerequisite, the AT_1R was shown to undergo a mechanically induced conformational change based when modeled after the bovine rhodopsin crystal structure (112), in which a dislocation and an anticlockwise rotation of transmembrane domain 7 toward the agonist-binding pocket occurred upon mechanical stimulation. Interestingly, mechanical membrane stretch stabilized a receptor conformation that was significantly different from that induced by agonist characterized by dislocation and rotation of transmembrane domain 6 out of the binding pocket, thereby veering away from transmembrane domain 3 (29, 45). The different agonist- or mechanically induced receptor conformations are displayed in Fig. 2. Furthermore, a distinct inactive receptor conformation induced by binding of an inverse agonist could be discerned.




Fig. 2. Agonist and mechanically induced receptor conformations. G protein-coupled receptors possess distinct receptor conformations: inactive (R), agonist-induced active ($R_{agonist}^*$) and mechanically induced active (R_{force}^*) receptor conformations. Active receptor conformations are characterized by conformational changes in transmembrane domains (displayed in red) resulting in G protein activation and activation of signal transduction pathways. It is still elusive whether mechanical GPCR activation requires interaction with the cytoskeleton or with the extracellular matrix (EC matrix) as proposed by the tethered model or whether the membrane model is applicable (indicated as membrane deformation). Agonist is shown in blue, and membrane stretch is indicated by two blue arrows.

The recent successful crystallization of the β_2 -adrenoceptor in its active conformation together with G_s (82) may serve as a better matrix for future receptor models. In recent years it became obvious that AT_1R and most probably also other GPCRs possess more than one active or inactive receptor conformation in analogy to the different states of ion channels. In line with this notion, different partial agonists were shown to induce several distinct active receptor conformations (116). In the performance of bioluminescence resonance energy transfer (BRET) experiments, it was demonstrated that membrane stretch favors an active AT_1R conformation that is recognized by β -arrestin (62), an observation that was recently verified by Rakesh et al. (81) in performing intermolecular fluorescence resonance energy transfer (FRET) experiments. In the latter study, G protein activation was not essential for mechanically induced β -arrestin recruitment. Of note, this study suggests the occurrence of a distinct β -arrestin conformation upon mechanical receptor activation, which is different from the one induced by agonist.

Although distinct mechanically and agonist-induced active receptor conformations could be discriminated, it still remains elusive whether mechanical receptor activation impacts agonist affinity. With a consideration that mechanical stimulation causes an increased sensitivity of the AT_1R toward agonist stimulation illustrated by a leftward shift of the concentration response curve (62), there is reason to assume that membrane stretch does at least not reduce agonist affinity. However, this issue still has to be finally resolved in the future. Moreover, it is not known whether GPCR agonists engage exactly the same signal transduction pathways as mechanical force.

Recently, Anfinogenova and colleagues (3) further analyzed the synergism between agonist- and mechanically induced receptor stimulation by measuring TRPC current activation in vascular smooth muscle cells as well as myogenic vasoconstriction. Surprisingly, after agonist stimulation with P2Y or thromboxane receptor agonists, subsequent mechanical stimulation in the presence of agonists did not enhance channel

activity or vasoconstriction, suggesting the absence of a functional synergism between these stimuli. However, it should be kept in mind that pretreatment with agonists inevitably entails receptor desensitization, resulting in a decreased secondary cell response. Furthermore, mechanosensitivity most probably depends on receptor density (62), and it is not clear whether P2Y and thromboxane receptor levels are higher in small resistance than in conduit arteries.

During the last years, some additional $G_{q/11}$ protein-coupled receptors have been demonstrated to be mechanosensitive. First of all, the $G_{q/11}$ protein-coupled bradykinin B_2 receptor exhibited mechanically induced conformational changes shown by intramolecular FRET (12). However, this study leaves open the question whether mechanical bradykinin B_2 receptor stimulation entails productive coupling to G proteins. Nevertheless, the mechanosensitivity of the B_2 bradykinin receptor may play a role in the sensing of fluid shear stress in endothelial cells.

Interestingly, many additional GPCRs were tested for their propensity to be activated by mechanical stimuli (for an over-

view of mechanosensitive GPCRs, see Table 1). Until now, very little is known about the mechanosensitivity of $G_{i/o}$ -coupled receptors, but first evidence suggests that fluid shear stress can activate $G_{i/o}$ -coupled formyl peptide-1 receptors as monitored by intermolecular FRET between G_i protein subunits (57). Although G_s -coupled β_2 -adrenoceptors did not respond to mechanical stimuli (62, 115), there is growing evidence that other G_s -coupled receptors might be inherently mechanosensitive. Recently, dopamine D_5 receptors in primary cilia were described as mechanosensors (1). Furthermore, in bone cells the parathyroid hormone type-1 receptor coupling to both G_s and $G_{q/11}$ proteins was shown to be activated by fluid shear stress (114). With an intramolecular FRET approach, fluid shear stress induced a distinct conformational change of the parathyroid hormone type-1 receptor. Interestingly, increasing the fluidity of the membrane using benzyl alcohol or cholesterol extraction, two maneuvers known to affect mechanosensitive ion channels (21), also led to conformational changes monitored by increased FRET levels. The observation that changes of the lipid composition of the cell membrane

Table 1. *Mechanosensitive GPCRs*

Receptor	Coupling	Expression System	Stimulation	Experimental Approach	Possible Function	Reference
AT ₁ R	$G_{q/11}$	Overexpression in HEK293 and in A7r5 cells, endogenous in vascular smooth muscle cells	Hypotonicity, direct membrane stretch, increased intravascular pressure	Whole cell measurements, Ca^{2+} imaging, inositol phosphate detection, BRET, measurements of myogenic vasoconstriction	Mechanosensor mediating myogenic vasoconstriction	(62)
		Overexpression in HEK293 and COS-7 cells, endogenous in cardiomyocytes	Direct membrane stretch, pressure overload	ERK phosphorylation, inositol phosphate detection, transthoracic echocardiography, heart morphometry, hemodynamic measurements	Pathophysiological role as mechanosensor inducing cardiac hypertrophy	(115)
		Overexpression of mutant receptors in HEK293 cells	Direct membrane stretch	Inhibition of angiotensin II binding, substituted cysteine accessibility mapping		(112)
		Overexpression in HEK293 cells, endogenous in cardiomyocytes	Hypotonicity, direct membrane stretch, pressure overload	ERK phosphorylation, intermolecular FRET, BRET	Pathophysiological role as mechanosensor inducing cardiac hypertrophy	(81)
B_2 R	$G_{q/11}$	Overexpression in bovine aortic endothelial cells	Fluid shear stress, hypotonicity	Intramolecular receptor FRET	Sensors of fluid shear stress in endothelial cells	(12)
D_5 R	G_s	Vascular endothelial cells	Fluid shear stress	Ca^{2+} measurements	Sensors of fluid shear stress in endothelial cells	(1)
ET _A R	$G_{q/11}$ / G_s	Overexpression in HEK293 cells	Hypotonicity	Whole cell measurements		(62)
FPR ₁	$G_{i/o}$	Endogenous in HL60 cells	Fluid shear stress	Intermolecular FRET between G_i protein subunits	Retraction of pseudopods in neutrophils	(57)
H ₁ R	$G_{q/11}$	Overexpression in HEK293 cells	Hypotonicity, direct membrane stretch	Whole cell and cell-attached patch-clamp measurements, Ca^{2+} imaging		(62)
M ₅ R	$G_{q/11}$	Overexpression in HEK293 cells	Hypotonicity, direct membrane stretch	Whole cell and cell-attached patch-clamp measurements		(62)
PTH ₁ R	G_s / $G_{q/11}$	Overexpression in HEK293 and in murine preosteoblastic cells	Fluid shear stress	Intramolecular receptor FRET	Bone growth	(114)
V _{1A} R	$G_{q/11}$	Overexpression in HEK293 cells	Hypotonicity	Whole cell measurements		(62)

Mechanosensitive G protein-coupled receptors (GPCRs). Overview of mechanosensitive GPCRs described so far, main signal transduction pathways, expression systems, stimuli and experimental approaches as well as proposed physiological or pathophysiological impact. See main text for definitions of abbreviations.

profoundly affect the activity of integral membrane proteins lends further credence to the idea of an activation model relying on changes of the lateral pressure profile.

Because GPCRs not only respond to mechanical or chemical stimuli but also sense alterations of the transmembrane potential (6), these membrane proteins appear to be endowed with polymodal-sensing capabilities to integrate a diverse array of different stimuli. This polymodal activation mechanism raises the intriguing question of whether the mechanosensitivity of GPCRs might be a general phenomenon that occurs in other mechanosensitive cells, tissues, and organs. This needs to be addressed in future studies. In addition to cardiac hypertrophy, mechanosensation and -transduction might also play key roles in other pathophysiological scenarios such as arteriosclerosis, glaucoma, chronic obstructive pulmonary disease and in the case of increased intracranial pressure.

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the author(s).

AUTHOR CONTRIBUTIONS

U.S., M.M.y.S., and T.G. drafted manuscript; U.S., M.M.y.S., and T.G. edited and revised manuscript; U.S., M.M.y.S., and T.G. approved final version of manuscript; M.M.y.S. prepared figures.

REFERENCES

- Abdul-Majeed S, Nauli SM. Dopamine receptor type 5 in the primary cilia has dual chemo- and mechano-sensory roles. *Hypertension* 58: 325–331, 2011.
- Alloui A, Zimmermann K, Mamet J, Duprat F, Noel J, Chemin J, Guy N, Blondeau N, Voilley N, Rubat-Coudert C, Borsotto M, Romey G, Heurteaux C, Reeh P, Eschalier A, Lazdunski M. TREK-1, a K⁺ channel involved in polymodal pain perception. *EMBO J* 25: 2368–2376, 2006.
- Anfinogenova Y, Brett SE, Walsh MP, Harraz OF, Welsh DG. Do TRPC-like currents and G protein-coupled receptors interact to facilitate myogenic tone development? *Am J Physiol Heart Circ Physiol* 301: H1378–H1388, 2011.
- Awayda MS, Ismailov II, Berdiev BK, Benos DJ. A cloned renal epithelial Na⁺ channel protein displays stretch activation in planar lipid bilayers. *Am J Physiol Cell Physiol* 268: C1450–C1459, 1995.
- Bae C, Sachs F, Gottlieb PA. The mechanosensitive ion channel Piezo1 is inhibited by the peptide GsMTx4. *Biochemistry* 50: 6295–6300, 2011.
- Ben-Chaim Y, Chanda B, Dascal N, Bezanilla F, Parnas I, Parnas H. Movement of ‘gating charge’ is coupled to ligand binding in a G-protein-coupled receptor. *Nature* 444: 106–109, 2006.
- Bessac BF, Fleig A. TRPM7 channel is sensitive to osmotic gradients in human kidney cells. *J Physiol* 582: 1073–1086, 2007.
- Bode F, Sachs F, Franz MR. Tarantula peptide inhibits atrial fibrillation. *Nature* 409: 35–36, 2001.
- Bounoutas A, Chalfie M. Touch sensitivity in *Caenorhabditis elegans*. *Pflügers Arch* 454: 691–702, 2007.
- Cantor RS. Lateral pressures in cell membranes: a mechanism for modulation of protein function. *J Phys Chem B* 101: 1723–1725, 1997.
- Carattino MD, Sheng S, Kleymann TR. Epithelial Na⁺ channels are activated by laminar shear stress. *J Biol Chem* 279: 4120–4126, 2004.
- Chachisvilis M, Zhang YL, Frangos JA. G protein-coupled receptors sense fluid shear stress in endothelial cells. *Proc Natl Acad Sci USA* 103: 15463–15468, 2006.
- Chalfie M. Neurosensory mechanotransduction. *Nat Rev Mol Cell Biol* 10: 44–52, 2009.
- Christensen AP, Corey DP. TRP channels in mechanosensation: direct or indirect activation? *Nat Rev Neurosci* 8: 510–521, 2007.
- Cipolla MJ, Gokina NI, Osol G. Pressure-induced actin polymerization in vascular smooth muscle as a mechanism underlying myogenic behavior. *FASEB J* 16: 72–76, 2002.
- Cipolla MJ, Osol G. Vascular smooth muscle actin cytoskeleton in cerebral artery forced dilatation. *Stroke* 29: 1223–1228, 1998.
- Clapham DE. TRP channels as cellular sensors. *Nature* 426: 517–524, 2003.
- Coste B, Mathur J, Schmidt M, Earley TJ, Ranade S, Petrus MJ, Dubin AE, Patapoutian A. Piezo1 and Piezo2 are essential components of distinct mechanically activated cation channels. *Science* 330: 55–60, 2010.
- D’Angelo G, Mogford JE, Davis GE, Davis MJ, Meininger GA. Integrin-mediated reduction in vascular smooth muscle [Ca²⁺]_i induced by RGD-containing peptide. *Am J Physiol Heart Circ Physiol* 272: H2065–H2070, 1997.
- Davis MJ, Hill MA. Signaling mechanisms underlying the vascular myogenic response. *Physiol Rev* 79: 387–423, 1999.
- Dedman A, Sharif-Naeini R, Folgering JH, Duprat F, Patel A, Honore E. The mechano-gated K(2P) channel TREK-1. *Eur Biophys J* 38: 293–303, 2009.
- Dietrich A, Kalwa H, Storch U, Mederos y Schnitzler M, Salanova B, Pinkenburg O, Dubrovskaja G, Essin K, Gollasch M, Birnbaumer L, Gudermann T. Pressure-induced and store-operated cation influx in vascular smooth muscle cells is independent of TRPC1. *Pflügers Arch* 455: 465–477, 2007.
- Doughty JM, Langton PD. Measurement of chloride flux associated with the myogenic response in rat cerebral arteries. *J Physiol* 534: 753–761, 2001.
- Drummond HA, Gebremedhin D, Harder DR. Degenerin/epithelial Na⁺ channel proteins: components of a vascular mechanosensor. *Hypertension* 44: 643–648, 2004.
- Drummond HA, Grifoni SC, Jernigan NL. A new trick for an old dogma: ENaC proteins as mechanotransducers in vascular smooth muscle. *Physiology (Bethesda)* 23: 23–31, 2008.
- Earley S, Straub SV, Brayden JE. Protein kinase C regulates vascular myogenic tone through activation of TRPM4. *Am J Physiol Heart Circ Physiol* 292: H2613–H2622, 2007.
- Earley S, Waldron BJ, Brayden JE. Critical role for transient receptor potential channel TRPM4 in myogenic constriction of cerebral arteries. *Circ Res* 95: 922–929, 2004.
- Emtage L, Gu G, Hartweg E, Chalfie M. Extracellular proteins organize the mechanosensory channel complex in *C. elegans* touch receptor neurons. *Neuron* 44: 795–807, 2004.
- Farrens DL, Altenbach C, Yang K, Hubbell WL, Khorana HG. Requirement of rigid-body motion of transmembrane helices for light activation of rhodopsin. *Science* 274: 768–770, 1996.
- Felsenfeld DP, Schwartzberg PL, Venegas A, Tse R, Sheetz MP. Selective regulation of integrin-cytoskeleton interactions by the tyrosine kinase Src. *Nat Cell Biol* 1: 200–206, 1999.
- Flavahan NA, Bailey SR, Flavahan WA, Mitra S, Flavahan S. Imaging remodeling of the actin cytoskeleton in vascular smooth muscle cells after mechanosensitive arteriolar constriction. *Am J Physiol Heart Circ Physiol* 288: H660–H669, 2005.
- Gabella G. Asymmetric distribution of dense bands in muscle cells of mammalian arterioles. *J Ultrastruct Res* 84: 24–33, 1983.
- Garry A, Fromy B, Blondeau N, Henrion D, Brau F, Gounon P, Guy N, Heurteaux C, Lazdunski M, Saumet JL. Altered acetylcholine, bradykinin and cutaneous pressure-induced vasodilation in mice lacking the TREK1 potassium channel: the endothelial link. *EMBO Rep* 8: 354–359, 2007.
- Geffeney SL, Cueva JG, Glauser DA, Doll JC, Lee TH, Montoya M, Karania S, Garakani AM, Pruitt BL, Goodman MB. DEG/ENaC but not TRP channels are the major mechano-electrical transduction channels in a *C. elegans* nociceptor. *Neuron* 71: 845–857, 2011.
- Gottlieb P, Folgering J, Maroto R, Raso A, Wood TG, Kurosky A, Bowman C, Bichet D, Patel A, Sachs F, Martinac B, Hamill OP, Honore E. Revisiting TRPC1 and TRPC6 mechanosensitivity. *Pflügers Arch* 455: 1097–1103, 2008.
- Greenwood IA, Large WA. Properties of a Cl⁻ current activated by cell swelling in rabbit portal vein vascular smooth muscle cells. *Am J Physiol Heart Circ Physiol* 275: H1524–H1532, 1998.
- Hamill OP. Twenty odd years of stretch-sensitive channels. *Pflügers Arch* 453: 333–351, 2006.
- Hein TW, Platts SH, Waitkus-Edwards KR, Kuo L, Mousa SA, Meininger GA. Integrin-binding peptides containing RGD produce coronary arteriolar dilation via cyclooxygenase activation. *Am J Physiol Heart Circ Physiol* 281: H2378–H2384, 2001.
- Ingber DE. Cellular mechanotransduction: putting all the pieces together again. *FASEB J* 20: 811–827, 2006.

40. Ingber DE. Mechanosensation through integrins: cells act locally but think globally. *Proc Natl Acad Sci USA* 100: 1472–1474, 2003.
41. Inoue R, Jensen LJ, Jian Z, Shi J, Hai L, Lurie AI, Henriksen FH, Salomonsson M, Morita H, Kawarabayashi Y, Mori M, Mori Y, Ito Y. Synergistic activation of vascular TRPC6 channel by receptor and mechanical stimulation via phospholipase C/diacylglycerol and phospholipase A2/omega-hydroxylase/20-HETE pathways. *Circ Res* 104: 1399–1409, 2009.
42. Inoue R, Jian Z, Kawarabayashi Y. Mechanosensitive TRP channels in cardiovascular pathophysiology. *Pharmacol Ther* 123: 371–385, 2009.
43. Jarajapu YP, Knot HJ. Role of phospholipase C in development of myogenic tone in rat posterior cerebral arteries. *Am J Physiol Heart Circ Physiol* 283: H2234–H2238, 2002.
44. Jin D, Takai S, Yamada M, Sakaguchi M, Miyazaki M. The functional ratio of chymase and angiotensin converting enzyme in angiotensin I-induced vascular contraction in monkeys, dogs and rats. *Jpn J Pharmacol* 84: 449–454, 2000.
45. Karnik SS, Gogonea C, Patil S, Saad Y, Takezako T. Activation of G-protein-coupled receptors: a common molecular mechanism. *Trends Endocrinol Metab* 14: 431–437, 2003.
46. Kato H, Osajima A, Uezono Y, Okazaki M, Tsuda Y, Tanaka H, Oishi Y, Izumi F, Nakashima Y. Involvement of PDGF in pressure-induced mesangial cell proliferation through PKC and tyrosine kinase pathways. *Am J Physiol Renal Physiol* 277: F105–F112, 1999.
47. Kirchenbuechler D, Born S, Kirchgessner N, Houben S, Hoffmann B, Merkel R. Substrate, focal adhesions, and actin filaments: a mechanical unit with a weak spot for mechanosensitive proteins. *J Phys Condens Matter* 22: 194109, 2010.
48. Kottgen M, Buchholz B, Garcia-Gonzalez MA, Kotsis F, Fu X, Doerken M, Boehlke C, Steffl D, Tauber R, Wegierski T, Nitschke R, Suzuki M, Kramer-Zucker A, Germino GG, Watnick T, Prenen J, Nilius B, Kuehn EW, Walz G. TRPP2 and TRPV4 form a polymodal sensory channel complex. *J Cell Biol* 182: 437–447, 2008.
49. Kung C. A possible unifying principle for mechanosensation. *Nature* 436: 647–654, 2005.
50. Lacolley P, Challande P, Boumaza S, Cohuet G, Laurent S, Boutouyrie P, Grimaud JA, Paulin D, Lamaziere JM, Li Z. Mechanical properties and structure of carotid arteries in mice lacking desmin. *Cardiovasc Res* 51: 178–187, 2001.
51. Laitko U, Juranka PF, Morris CE. Membrane stretch slows the concerted step prior to opening in a Kv channel. *J Gen Physiol* 127: 687–701, 2006.
52. Laitko U, Morris CE. Membrane tension accelerates rate-limiting voltage-dependent activation and slow inactivation steps in a Shaker channel. *J Gen Physiol* 123: 135–154, 2004.
53. Lehtonen JY, Kinnunen PK. Phospholipase A₂ as a mechanosensor. *Biophys J* 68: 1888–1894, 1995.
54. Levick JR. *An Introduction to Cardiovascular Physiology*. London: Hodder Arnold, 2010, p. 222, 242–243, 250–251.
55. Loufrani L, Matrougui K, Li Z, Levy BI, Lacolley P, Paulin D, Henrion D. Selective microvascular dysfunction in mice lacking the gene encoding for desmin. *FASEB J* 16: 117–119, 2002.
56. Maingret F, Patel AJ, Lesage F, Lazdunski M, Honore E. Mechano- or acid stimulation, two interactive modes of activation of the TREK-1 potassium channel. *J Biol Chem* 274: 26691–26696, 1999.
57. Makino A, Prossnitz ER, Bunemann M, Wang JM, Yao W, Schmid-Schonbein GW. G protein-coupled receptors serve as mechanosensors for fluid shear stress in neutrophils. *Am J Physiol Cell Physiol* 290: C1633–C1639, 2006.
58. Maroto R, Raso A, Wood TG, Kurosky A, Martinac B, Hamill OP. TRPC1 forms the stretch-activated cation channel in vertebrate cells. *Nat Cell Biol* 7: 179–185, 2005.
59. Martinez-Lemus LA, Crow T, Davis MJ, Meininger GA. $\alpha_5\beta_3$ - and $\alpha_5\beta_1$ -integrin blockade inhibits myogenic constriction of skeletal muscle resistance arterioles. *Am J Physiol Heart Circ Physiol* 289: H322–H329, 2005.
60. Matsumoto H, Baron CB, Coburn RF. Smooth muscle stretch-activated phospholipase C activity. *Am J Physiol Cell Physiol* 268: C458–C465, 1995.
61. Mederos y Schnitzler M, Storch U, Gudermann T. AT₁ receptors as mechanosensors. *Curr Opin Pharmacol* 11: 112–116, 2011.
62. Mederos y Schnitzler M, Storch U, Meibers S, Nurwakagari P, Breit A, Essin K, Gollasch M, Gudermann T. Gq-coupled receptors as mechanosensors mediating myogenic vasoconstriction. *EMBO J* 27: 3092–3103, 2008.
63. Miyazaki M, Takai S. Local angiotensin II-generating system in vascular tissues: the roles of chymase. *Hypertens Res* 24: 189–193, 2001.
64. Mogford JE, Davis GE, Platts SH, Meininger GA. Vascular smooth muscle $\alpha_v\beta_3$ integrin mediates arteriolar vasodilation in response to RGD peptides. *Circ Res* 79: 821–826, 1996.
65. Montell C, Birnbaumer L, Flockerzi V. The TRP channels, a remarkably functional family. *Cell* 108: 595–598, 2002.
66. Morita H, Honda A, Inoue R, Ito Y, Abe K, Nelson MT, Brayden JE. Membrane stretch-induced activation of a TRPM4-like nonselective cation channel in cerebral artery myocytes. *J Pharm Sci* 103: 417–426, 2007.
67. Morris CE. Voltage-gated channel mechanosensitivity: fact or friction? *Front Physiol* 2: 25, 2011.
68. Muraki K, Iwata Y, Katanosaka Y, Ito T, Ohya S, Shigekawa M, Imaizumi Y. TRPV2 is a component of osmotically sensitive cation channels in murine aortic myocytes. *Circ Res* 93: 829–838, 2003.
69. Na S, Trache A, Trzeciakowski J, Sun Z, Meininger GA, Humphrey JD. Time-dependent changes in smooth muscle cell stiffness and focal adhesion area in response to cyclic equibiaxial stretch. *Ann Biomed Eng* 36: 369–380, 2008.
70. Narayanan J, Imig M, Roman RJ, Harder DR. Pressurization of isolated renal arteries increases inositol trisphosphate and diacylglycerol. *Am J Physiol Heart Circ Physiol* 266: H1840–H1845, 1994.
71. Nauli SM, Alenghat FJ, Luo Y, Williams E, Vassilev P, Li X, Elia AE, Lu W, Brown EM, Quinn SJ, Ingber DE, Zhou J. Polycystins 1 and 2 mediate mechanosensation in the primary cilium of kidney cells. *Nat Genet* 33: 129–137, 2003.
72. Nelson MT, Conway MA, Knot HJ, Brayden JE. Chloride channel blockers inhibit myogenic tone in rat cerebral arteries. *J Physiol* 502: 259–264, 1997.
73. Nilius B, Prenen J, Tang J, Wang C, Owsianik G, Janssens A, Voets T, Zhu MX. Regulation of the Ca²⁺ sensitivity of the nonselective cation channel TRPM4. *J Biol Chem* 280: 6423–6433, 2005.
74. Numata T, Shimizu T, Okada Y. Direct mechano-stress sensitivity of TRPM7 channel. *Cell Physiol Biochem* 19: 1–8, 2007.
75. Numata T, Shimizu T, Okada Y. TRPM7 is a stretch- and swelling-activated cation channel involved in volume regulation in human epithelial cells. *Am J Physiol Cell Physiol* 292: C460–C467, 2007.
76. Orr AW, Helmke BP, Blackman BR, Schwartz MA. Mechanisms of mechanotransduction. *Dev Cell* 10: 11–20, 2006.
77. Osol G, Brekke JF, McElroy-Yaggy K, Gokina NI. Myogenic tone, reactivity, and forced dilatation: a three-phase model of in vitro arterial myogenic behavior. *Am J Physiol Heart Circ Physiol* 283: H2260–H2267, 2002.
78. Osol G, Laher I, Kelley M. Myogenic tone is coupled to phospholipase C and G protein activation in small cerebral arteries. *Am J Physiol Heart Circ Physiol* 265: H415–H420, 1993.
79. Patel AJ, Honore E, Maingret F, Lesage F, Fink M, Duprat F, Lazdunski M. A mammalian two pore domain mechano-gated S-like K⁺ channel. *EMBO J* 17: 4283–4290, 1998.
80. Phillips R, Ursell T, Wiggins P, Sens P. Emerging roles for lipids in shaping membrane-protein function. *Nature* 459: 379–385, 2009.
81. Rakesh K, Yoo B, Kim IM, Salazar N, Kim KS, Rockman HA. beta-Arestin-biased agonism of the angiotensin receptor induced by mechanical stress. *Sci Signal* 3: ra46, 2010.
82. Rasmussen SG, DeVree BT, Zou Y, Kruse AC, Chung KY, Kobilka TS, Thian FS, Chae PS, Pardon E, Calinski D, Mathiesen JM, Shah ST, Lyons JA, Caffrey M, Gellman SH, Steyaert J, Skiniotis G, Weis WI, Sunahara RK, Kobilka BK. Crystal structure of the beta2 adrenergic receptor-Gs protein complex. *Nature* 477: 549–555, 2011.
83. Sadoshima J, Jahn L, Takahashi T, Kulik TJ, Izumo S. Molecular characterization of the stretch-induced adaptation of cultured cardiac cells. An in vitro model of load-induced cardiac hypertrophy. *J Biol Chem* 267: 10551–10560, 1992.
84. Sawada Y, Tamada M, Dubin-Thaler BJ, Cherniavskaya O, Sakai R, Tanaka S, Sheetz MP. Force sensing by mechanical extension of the Src family kinase substrate p130Cas. *Cell* 127: 1015–1026, 2006.
85. Sharif-Naeini R, Folgering JH, Bichet D, Duprat F, Delmas P, Patel A, Honore E. Sensing pressure in the cardiovascular system: Gq-coupled mechanoreceptors and TRP channels. *J Mol Cell Cardiol* 48: 83–89, 2010.

86. Sharif-Naeini R, Folgering JH, Bichet D, Duprat F, Lauritzen I, Arhatte M, Jodar M, Dedman A, Chatelain FC, Schulte U, Retailleau K, Loufrani L, Patel A, Sachs F, Delmas P, Peters DJ, Honore E. Polycystin-1 and -2 dosage regulates pressure sensing. *Cell* 139: 587–596, 2009.
87. Shi S, Ghosh DD, Okumura S, Carattino MD, Kashlan OB, Sheng S, Kleyman TR. Base of the thumb domain modulates epithelial sodium channel gating. *J Biol Chem* 286: 14753–14761, 2011.
88. Small JV. Geometry of actin-membrane attachments in the smooth muscle cell: the localisations of vinculin and alpha-actinin. *EMBO J* 4: 45–49, 1985.
89. Spassova MA, Hewavitharana T, Xu W, Soboloff J, Gill DL. A common mechanism underlies stretch activation and receptor activation of TRPC6 channels. *Proc Natl Acad Sci USA* 103: 16586–16591, 2006.
90. Suchyna TM, Tape SE, Koeppe RE 2nd, Andersen OS, Sachs F, Gottlieb PA. Bilayer-dependent inhibition of mechanosensitive channels by neuroactive peptide enantiomers. *Nature* 430: 235–240, 2004.
91. Sukharev SI, Martinac B, Arshavsky VY, Kung C. Two types of mechanosensitive channels in the *Escherichia coli* cell envelope: solubilization and functional reconstitution. *Biophys J* 65: 177–183, 1993.
92. Tang DD, Anfinogenova Y. Physiologic properties and regulation of the actin cytoskeleton in vascular smooth muscle. *J Cardiovasc Pharmacol Ther* 13: 130–140, 2008.
93. Tsiokas L, Arnould T, Zhu C, Kim E, Walz G, Sukhatme VP. Specific association of the gene product of PKD2 with the TRPC1 channel. *Proc Natl Acad Sci USA* 96: 3934–3939, 1999.
94. Vandenburg HH, Shansky J, Karlisch P, Solerssi RL. Mechanical stimulation of skeletal muscle generates lipid-related second messengers by phospholipase activation. *J Cell Physiol* 155: 63–71, 1993.
95. VanLandingham LG, Gannon KP, Drummond HA. Pressure-induced constriction is inhibited in a mouse model of reduced β ENaC. *Am J Physiol Regul Integr Comp Physiol* 297: R723–R728, 2009.
96. Vennekens R, Nilius B. Insights into TRPM4 function, regulation and physiological role. *Handb Exp Pharmacol* 179: 269–285, 2007.
97. Vilardaga JP, Bunemann M, Krasel C, Castro M, Lohse MJ. Measurement of the millisecond activation switch of G protein-coupled receptors in living cells. *Nat Biotechnol* 21: 807–812, 2003.
98. Voets T, Nilius B. TRPCs, GPCRs and the Bayliss effect. *EMBO J* 28: 4–5, 2009.
99. Vriens J, Owsianik G, Fisslthaler B, Suzuki M, Janssens A, Voets T, Morisseau C, Hammock BD, Fleming I, Busse R, Nilius B. Modulation of the Ca²⁺ permeable cation channel TRPV4 by cytochrome P450 epoxygenases in vascular endothelium. *Circ Res* 97: 908–915, 2005.
100. Waite M. Approaches to the study of mammalian cellular phospholipases. *J Lipid Res* 26: 1379–1388, 1985.
101. Waitkus-Edwards KR, Martinez-Lemus LA, Wu X, Trzeciakowski JP, Davis MJ, Davis GE, Meininger GA. α (4) β (1) Integrin activation of L-type calcium channels in vascular smooth muscle causes arteriole vasoconstriction. *Circ Res* 90: 473–480, 2002.
102. Wang S, Meng F, Mohan S, Champaneri B, Gu Y. Functional ENaC channels expressed in endothelial cells: a new candidate for mediating shear force. *Microcirculation* 16: 276–287, 2009.
103. Wang Y, Botvinick EL, Zhao Y, Berns MW, Usami S, Tsien RY, Chien S. Visualizing the mechanical activation of Src. *Nature* 434: 1040–1045, 2005.
104. Watanabe H, Vriens J, Prenen J, Droogmans G, Voets T, Nilius B. Anandamide and arachidonic acid use epoxyeicosatrienoic acids to activate TRPV4 channels. *Nature* 424: 434–438, 2003.
105. Welsh DG, Morielli AD, Nelson MT, Brayden JE. Transient receptor potential channels regulate myogenic tone of resistance arteries. *Circ Res* 90: 248–250, 2002.
106. Welsh DG, Nelson MT, Eckman DM, Brayden JE. Swelling-activated cation channels mediate depolarization of rat cerebrovascular smooth muscle by hyposmolarity and intravascular pressure. *J Physiol* 527: 139–148, 2000.
107. Wijetunge S, Hughes AD. Src family tyrosine kinases mediate contraction of rat isolated tail arteries in response to a hyposmotic stimulus. *J Hypertens* 25: 1871–1878, 2007.
108. Wu X, Davis GE, Meininger GA, Wilson E, Davis MJ. Regulation of the L-type calcium channel by α 5 β 1 integrin requires signaling between focal adhesion proteins. *J Biol Chem* 276: 30285–30292, 2001.
109. Wu X, Yang Y, Gui P, Sohma Y, Meininger GA, Davis GE, Braun AP, Davis MJ. Potentiation of large conductance, Ca²⁺-activated K⁺ (BK) channels by α 5 β 1 integrin activation in arteriolar smooth muscle. *J Physiol* 586: 1699–1713, 2008.
110. Yamazaki J, Duan D, Janiak R, Kuenzli K, Horowitz B, Hume JR. Functional and molecular expression of volume-regulated chloride channels in canine vascular smooth muscle cells. *J Physiol* 507: 729–736, 1998.
111. Yasuda N, Akazawa H, Qin Y, Zou Y, Komuro I. A novel mechanism of mechanical stress-induced angiotensin II type 1-receptor activation without the involvement of angiotensin II. *Naunyn Schmiedebergs Arch Pharmacol* 377: 393–399, 2008.
112. Yasuda N, Miura S, Akazawa H, Tanaka T, Qin Y, Kiya Y, Imaizumi S, Fujino M, Ito K, Zou Y, Fukuhara S, Kunimoto S, Fukuzaki K, Sato T, Ge J, Mochizuki N, Nakaya H, Saku K, Komuro I. Conformational switch of angiotensin II type 1 receptor underlying mechanical stress-induced activation. *EMBO Rep* 9: 179–186, 2008.
113. Yin J, Kuebler WM. Mechanotransduction by TRP channels: general concepts and specific role in the vasculature. *Cell Biochem Biophys* 56: 1–18, 2009.
114. Zhang YL, Frangos JA, Chachisvilis M. Mechanical stimulus alters conformation of type 1 parathyroid hormone receptor in bone cells. *Am J Physiol Cell Physiol* 296: C1391–C1399, 2009.
115. Zou Y, Akazawa H, Qin Y, Sano M, Takano H, Minamino T, Makita N, Iwanaga K, Zhu W, Kudoh S, Toko H, Tamura K, Kihara M, Nagai T, Fukamizu A, Umemura S, Iiri T, Fujita T, Komuro I. Mechanical stress activates angiotensin II type 1 receptor without the involvement of angiotensin II. *Nat Cell Biol* 6: 499–506, 2004.
116. Zurn A, Zabel U, Vilardaga JP, Schindelin H, Lohse MJ, Hoffmann C. Fluorescence resonance energy transfer analysis of α 2a-adrenergic receptor activation reveals distinct agonist-specific conformational changes. *Mol Pharmacol* 75: 534–541, 2009.